

PCT/EP

98/08004

09/581847



Eur päisches  
Patentamt

European  
Patent Office

Office eur péen  
des brevets

REC'D 09 MAR 1999

WIPO

PCT

EP98/08004

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

97122216.1

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

DEN AAG, DEN  
THE GUE,  
A H, LE

19/02/99

B. Scholz



Eur päisches  
Patentamt

Eur pean  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.: 97122216.1  
Demande n°:

Anmeldetag:  
Date of filing: 17/12/97  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
MERCK PATENT GmbH  
64293 Darmstadt  
GERMANY  
Limerick Ltd  
Tel Aviv 64256

ISRAEL  
Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

Method and composition for combatting insects and viruses transmitted to plants thereby

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

/

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

M 17-10-97

3

Merck Patent Gesellschaft  
mit beschränkter Haftung  
64271 Darmstadt

**METHOD AND COMPOSITION FOR  
COMBATTING INSECTS AND VIRUSES  
TRANSMITTED TO PLANTS THEREBY**

## METHOD AND COMPOSITION FOR COMBATTING INSECTS AND VIRUSES TRANSMITTED TO PLANTS THEREBY

5 The present invention relates to a method and composition for combatting insects and insect-transmitted plant viruses by repelling insects from the plants.

10 It is a continuing objective of persons engaged in agriculture, to increase yields of crops such as vegetables, especially by combatting plant pests such as plant viruses introduced by insects. Use of insecticides by no means provides a complete answer to this problem, due to the known phenomenon of increasing insect resistance to insecticides.

15 An article by Smith, F.F. and Webb, R.E., entitled „Repelling Aphids by Reflective Surfaces ...“ at pages 631-639 of „Viruses, Vectors and Vegetation“ (Karl Maramorosch, ed.), Interscience Publishers, 1969, reviews earlier work describing the reaction aphids to different colors, and in particular the finding that aphids are attracted to yellow or modified yellow objects (such as a yellow pan), but are influenced also by the nature or color background of the background surrounding the yellow, e.g. unpainted aluminum pans surrounding yellow pans repels the aphids. The review also mentioned as a known fact, that sprayed foliage, especially that having a whitish residue, attracts aphids to land. The authors of this review then reported the results of their own experiments, in which it was found that foliage sprays of aluminum repelled as many aphids as aluminum sheets. The aluminum sprays, however, suffered from a number of disadvantages, namely, either the spray deposits adhered to plant for only 2-3 days and thereafter lost their effectiveness, or on rough hairy plants (such as cucumber, canteloupe or tobacco) the sprays left a dull gray deposit which did not repel aphids. More persistent aluminum sprays containing adhesive were found to stunt growth of foliage. Using aluminum sheets as mulches, in experiments with bush squashes, the authors found that maximum aphid repellency was obtained when 50% of the ground was thus covered, and that much higher yields were obtained than when using parathion spray, or in a control experiment. It was further reported

20  
25  
30  
35

that aluminum sheet material was not succesful in repelling aphids from all plant species.

5 JP 4190733A (published July, 1992) discloses a multi-film for repelling harmful insects and accelerating growth of vegetation, prepared from a composition containing thermoplastic resin, metal aluminium powder and mica, and optionally, e.g. antioxidants, and antistatic and UV absorbing agents. JP 4152833A (published May, 1992) discloses a similar product for reducing the effect of harmful insects on agricultural products, but in  
10 which (in particular) the mica of the previous composition is replaced by mica that is  $\text{TiO}_2$ -coated as pearl pigment.

The entire disclosure of the literature reference and the Japanese patent publications mentioned above is explicitly incorporated herein by  
15 reference.

Thus, the tendency of the prior art known to Applicant, so far as repelling insects by use of reflective materials is concerned, is to utilize sheet material which is either itself reflective as in the case of the aluminum  
20 sheet mulches described in the Smith and Webb articles, or which is apparently reflective by virtue of including reflective particles as in the case of the above-mentioned published JP application.

However, these sheet materials suffer two important disadvantages.  
25 Firstly, they tend to heat the microenvironment of the plants they are intended to protect, so that in warm climates the plants could be damaged or even destroyed by intense heat thus induced by these sheets. Secondly, these sheets are expensive, of the order of US \$500 per 0.1 hectare.

30 Marco, S. in Phytopathology, 76: 1344-1348 (1986) reported the results of experiments in which potato crops were sprayed weekly with whitewashes. It was found that potato leaf roll virus was reduced by 0-61% and potato virus Y by 0-68%; and also that a number of aphid species were repelled  
35 by whitewash while one species was attracted (as determined by comparing the number of aphids attracted to sprayed traps and control

traps). However, the use of whitewashes suffers serious disadvantages, namely: the repellant effect on insects is not powerful enough; increasing the concentration in the spray could harm the plants; and the sprayed crops would require special treatment to remove the whitewash from the fruit or vegetables in question. Whitewashes are in fact hardly used by farmers at all for crop protection, except for painting the bark of trees.

It has been surprisingly found in accordance with the present invention, that insects can be effectively repelled from growing plants in a manner which is considerably more economic than the prevalent method of using reflective mulching sheets, which avoids damage to or destruction of plants caused by use of such sheets in warm climates, and which avoids the above-noted disadvantage of whitewashes.

The present invention accordingly provides a method for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises applying to a locus or loci selected from surfaces of growing plants and their background, a composition comprising reflective particles of at least one substance, provided that this is not solely uncoated aluminum, whereby said composition is effective to repel said insects and thus protect the plants.

In another aspect, the invention provides a composition for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises reflective particles of at least one substance, together with at least one agriculturally acceptable diluent, carrier or adjuvant, provided that said substance is not solely uncoated aluminum.

The present invention provides a means for avoiding, at least to a large extent, the damaging effect on growing plants of destructive insects and insect-transmitted plant viruses. The method of the invention also reduces the temperature of the microenvironment of the plants, which can be of great importance in warm climates; e.g., leaf burns in zucchini are avoided, and the quality and sweetness of melons are improved.

5 The means of applying a composition comprising reflective particles to the surfaces of growing plants and/or their background in accordance with the invention may be any such means known in the art. Spraying is presently preferred, but this does not exclude, e.g., the use of powders for dusting plants.

10 As will be apparent from the preceding paragraph, the composition of the invention, which may be used in the method of the invention, are essentially admixtures which may take any form known per se in the art, e.g. they may be in the form of suspensions or dispersions for applying by means of a spray, or they may take the form of dusting powders. In any event, they may contain agriculturally acceptable conventional hydrophilic or lipophilic diluents, carriers or adjuvants. Examples for diluents are mineral oil, vegetable oil or water. For example, aqueous suspensions or  
15 dispersions may contain surface-active agents and/or adhesives for attaching the reflective particles to plants. Dusting powders may contain, e.g. solid carriers or diluents such as chalk or diatomaceous earth. The term „reflective particles“ is intended to include light-reflective powders, platelets, leaflets and laminae, and include light-reflective substances  
20 known for use in pigments as set forth in more detail infra.

As the reflective particles (or powders or platelets) which may be utilized in the practice of the method of the invention, there may be used, e.g., sheet silicates like talk, kaolin, sericit or mica, glass platelets,  $\text{SiO}_2$  flakes,  
25 ceramic flakes, platelet-shaped barium sulfate, platelet-shaped alumina or micaceous iron oxide, particularly when coated with layers of metal oxides like  $\text{TiO}_2$  and/or  $\text{Fe}_2\text{O}_3$ . Coated mica platelets are for example marketed under the Registered Trade Marks „Iriodin“ (Merck KGaA, Darmstadt, Germany) and „Mearlin“ or „MagnaPearl“ (The Mearl Corporation, New  
30 York, U.S.A.), for use as pearl luster pigments.

A particular non-limiting example of the Mearlin range of coated mica platelets, found to be useful in the present invention, is Mearlin Card Silver BN001, which contains generally 64-67% mica, 22-31%  $\text{TiO}_2$ , 2-4%  
35 graphite („Graphitan 6154“) and 0.1-0.5%  $\text{SnO}_2$ ; the usefulness of this product in the practice of the invention does not of course preclude the use

of  $\text{TiO}_2$ -coated mica wherein the coating contains different amounts of graphite and  $\text{SnO}_2$ , of which contains only one or neither of these ingredients, or which contains other minor ingredients. A further example of the reflective particles, powders or platelets which may be utilized in the practice of the invention, is constituted by  $\text{BiOCl}$  crystals marketed under the Registered Trade Mark „Mearlite“ (The Mearl Corporation).

As the reflective particles (or powders or platelets) which may be utilized in the practice of the invention, a further example consists of aluminum coated with epoxy and/or polyester, and marketed under the trade name „Debrex“ by Debra Incorporated (New York, U.S.A.). Reflective copper or bronze particles, etc., are also potentially useful in practising the invention.

Further examples of the reflective particles, powders or platelets which may be utilized in the practice of the method of the invention, are constituted by pigment-free iridescent glitter also marketed by Debra Incorporated, such as (i) „HT glitter“, (ii) „LR glitter“ and (iii) „SR glitter“. The essential polymers in these compositions appear to be: (i) polyethylene terephthalate + acrylates copolymer; (ii) polybutylene terephthalate + acrylates copolymer + ethylene/vinyl acetate copolymer; (iii) polyethylene terephthalate + polybutylene terephthalate + ethylene/vinyl acetate copolymer.

A person of the art will of course appreciate that in accordance with the invention, the active insect-repellent substances are selected for their light-reflective, pearlescent or glitter properties, and thus the actual chemical composition of the particles is a secondary consideration, except that since the present invention is principally concerned with treating crops intended for human consumption it will of course be highly desirable that the active insect-repellant substances be non-toxic. A person of the art will also be aware that where such reflective particles, powders or platelets are supplied by the manufacturer in a vehicle for use as pigments, such vehicles could make these products unsuitable for agricultural applications, where predominantly aqueous vehicles would normally be used. In other words, the active insect-repellant substances should normally be initially in solvent-free form and will then be introduced for utilitaion purposes into



aqueous medium; this general principle does not of course exclude the possibility of utilizing such a commercially available substance in a vehicle suitable for agricultural application, provided that toxic substances are absent therefrom.

5

Without prejudice to the generality of the invention, it is presently believed that the method of the invention will be especially applicable to repelling aphids, leafhoppers, *Lariomyza Bryoniae*, white flies and thrips, from growing plants, thus protecting them from these insects as well as from plant viruses transmitted by these insects.

10

The invention will now be illustrated by the following Examples.

#### Example 1

15

Six yellow plates (20 x 40 cm) were placed in a row at a distance of 15 meters from the edge of and along one side of a cotton field, and six such yellow plates were placed similarly in a second row, but at a distance of 25 meters from the edge of a field. The plates were sprayed (two from each row) with, respectively, 1% aqueous suspensions (containing 0.1% „Shatach 90“ surface active agent) of mica coated with titanium dioxide (M.T.O.) and mica coated with titanium oxide and graphite (BN001 as described above), the plates having been sanded previously to improve distribution of the spray (which was not however distributed uniformly).

20

Two control plates from each row were not sprayed. Each such spraying was followed by spraying with a transparent glue „Rimiput“ for adhering leaf-hoppers landing on the plates. The experiment was commenced in the evening of day 1 and the plates were collected in the afternoon of day 5, the number of leaf-hoppers on each plate being as follows:

25  
30

35

Run No.	Control	M.T.O.	BN001
1	446	34	33
2	598	45	28
3	408	45	21
4	534	40	35
Mean	496.5	41	29.9
Standard Deviation	74.32	4.53	5.4

Summary of Results. M.T.O. and BN001 were, 91.73 and 94% effective, respectively, in reducing the no. of captured leaf-hoppers, compared with control, while the BN001 reduced the no. of captured leaf-hoppers by 28.56%, compared with M.T.O.

#### Example 2

Twenty yellow plates (20 x 20 cm) were placed at random in a melon field, and were sprayed in batches of five with, respectively, 1% aqueous suspensions (containing 0.1% „Shatach 90“ surface active agent) of uncoated mica (M), mica coated with titanium oxide (M.T.O.) and mica coated with titanium oxide and graphite (BN001), the plates having been sanded previously to improve distribution of the spray (which was not however distributed uniformly). Five control plates were not sprayed. Each such spraying was followed by spraying with a transparent glue as in Example 1. After four days the number of white flies and leaf-hoppers on each plate was as follows:

(a) White flies Effect of spraying on the number of white flies collected:

Run No.	Control	M	M.T.O.	BN001
1	432	368	344	124
2	469	421	132	321
3	500	370	217	256
4	493	437	203	280
5	535	375	278	132
Mean	485.8	394.2	234.8	176.4
Standard Deviation	34.2	28.95	71.68	108.89

Summary of Results. M.T.O. and BN001 were approximately 50-60% effective in reducing the incidence of captured white flies, whereas M also reduced the incidence of captured white flies but by <20%.

(b) Leaf hoppers Effect of spraying on the number of leaf hoppers collected:

Run No.	Control	M	M.T.O.	BN001
1	39	97	6	5
2	69	48	5	10
3	55	28	3	2
4	92	83	15	2
5	77	43	3	5
Mean	66.4	56.8	6.4	4.8
Standard Deviation	18.19	25.9	4.5	2.9

Summary of Results. M.T.O. and BN001 were ~90% effective in reducing the incidence of captured leaf-hoppers; M also showed a (non-statistical) tendency to reduce the incidence of captured leaf-hoppers.

Example 3

On July 25, 1991, Galia melon was sown in an experimental plot containing four rows, approximately seventy plants per row. The plants were divided into three groups: the control group was unsprayed, one group was sprayed with a commercial oil („Virol“) and the other group was sprayed according to the present invention (this spray was an aqueous suspension of 1% epoxy-coated aluminum particles, containing also (approximately) 0.1% surface-active agent and 0.1% polvinyl agricultural adhesive). The sprayings were carried out after one leaf had formed (July 30), and again on August 4, 9, 15, 20, 25 and 30, and on September 11 and 16. The spray left only slight traces on the vegetation, which could be seen only with difficulty. Monitoring of Cucumber Mosaic Virus (CMV) (transmitted by aphids) was carried out on three dates prior to picking; monitoring of Cucumber Yellow Vein Clearing Virus (CYVV) (transmitted by white flies) was carried out only on the first date, since this virus had disappeared by the two later days. In each case monitoring was by observation, any doubtful cases being remitted for laboratory examination for serological confirmation of infection. The results of this experiment, in which each group for which data is given contained 62-75 plants, were as follows:

Percentages of infected plants in group

Date	Group	Control	„Virol“	Epoxy-coated A1
August 23 (CMV)	1	13.3	7.7	1.6
	2	10.0	12.5	4.4
	3	11.3	2.9	1.5
	4	7.7	7.4	1.4
Mean		10.7	7.7	2.3
August 30 (CMV)	1	20.0	9.2	1.6
	2	12.9	18.1	4.4
	3	16.1	4.3	1.5
	4	7.7	7.4	1.4
Mean		14.3	9.9	2.3

5	Sept. 16	1	20.0	9.2	1.6
	(CMV)	2	12.9	18.1	4.4
		3	16.1	10.1	4.6
		4	23.1	8.8	2.9
		Mean	18.0	11.7	3.4

10	Date	Group	Control	„Virol“	Epoxy-coated A1
	August 23	1	24.0	20.0	6.3
	(CYVV)	2	21.4	5.6	1.5
		3	6.5	21.7	1.5
		4	7.7	2.9	10.1
		Mean	15.4	12.4	4.9

15     Conclusions. The above-tabulated results show that the rate of infection in the oil-treated plants was moderately lower than the control, while the plants sprayed in accordance with the invention showed a rate of infection considerably lower than the control.

20     While particular embodiments of the invention have been described above, the skilled person will be aware that many variation and modifications may be made. The present invention includes all such variations and modifications which lie within the spirit, scope or concept of the present invention, and which would be apparent to the skilled addressee, on

25     reading the present disclosure and the claims which follow.

30

35

## CLAIMS

1. A method for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises applying to a locus or loci selected from surfaces of growing plants and their background, a composition comprising reflective particles of at least one substance, provided that this is not solely uncoated aluminum, whereby said composition is effective to repel said insects and thus protect the plants.
2. Method according to claim 1, wherein said composition is adapted for applying to said surfaces in the form of a spray or dusting powder.
3. Method according to claim 2, wherein said composition is an aqueous suspension or dispersion of said reflective particles.
4. Method according to claim 3, wherein said aqueous suspension or dispersion comprises additionally at least one surface active agent or adhesive.
5. Method according to claim 1, wherein said particles are selected from the group consisting of mica, mica coated with  $\text{TiO}_2$ , mica coated with  $\text{Fe}_2\text{O}_3$ , mica coated with both  $\text{TiO}_2$  and  $\text{Fe}_2\text{O}_3$ , mica coated with both  $\text{TiO}_2$  and graphite, reflective copper, reflective bronze, reflective coated aluminum, and  $\text{BiOCl}$ .
6. Method according to claim 5, wherein said mica coated with  $\text{TiO}_2$  also contains in the coating at least one of graphite and  $\text{SnO}_2$ .
7. Method according to claim 1, wherein said particles are constituted by pigment-free iridescent glitter.
8. Method according to claim 1, wherein said particles comprise pearlescent particles.

9. Method according to claim 1, wherein said insects are selected from aphids, leafhoppers, Lariomyza Bryoniae, white flies and thrips.
- 5 10. A composition for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises reflective particles of at least one substance, together with at least one agriculturally acceptable diluent, carrier or adjuvant, provided that said substance is not solely uncoated aluminum.
- 10 11. A composition according to claim 10, which is adapted for application to the surfaces of growing plants in the form of a spray or dusting powder.
- 15 12. A composition according to claim 11, which is an aqueous suspension or dispersion of said reflective particles.
- 20 13. A composition according to claim 12, wherein said aqueous suspension or dispersion comprises additionally at least one surface active agent or adhesive.
- 25 14. A composition according to claim 10, wherein said particles are selected from the group consisting of mica, mica coated with  $\text{TiO}_2$ , mica coated with  $\text{Fe}_2\text{O}_3$ , mica coated with both  $\text{TiO}_2$  and  $\text{Fe}_2\text{O}_3$ , mica coated with both  $\text{TiO}_2$  and graphite, reflective copper, reflective bronze, reflective aluminum, and  $\text{BiOCl}$ .
- 30 15. A composition according to claim 14, wherein said mica coated with  $\text{TiO}_2$  also contains in the coating at least one of graphite and  $\text{SnO}_2$ .
- 35 16. A composition according to claim 10, wherein said particles are constituted by pigment-free iridescent glitter.
17. A composition according to claim 10, wherein said particles comprise pearlescent particles.

### Conclusion

5 The invention relates to a method for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises applying to a locus or loci selected from surfaces of growing plants and their background, a composition comprising reflective particles of at least one substance, provided that this is not solely uncoated aluminum, whereby said composition is effective to repel said insects and thus protect the plants.

10 Furthermore the invention relates to a composition for protecting growing plants from insects and from insect-transmitted plant viruses, which comprises reflective particles of at least one substance, together with at least one agriculturally acceptable diluent, carrier or adjuvant, provided  
15 that said substance is not solely uncoated aluminum.

20

25

30

35